

Supplement to “Nonparametric Threshold Detection for Cost Distributions”

Benedikt Funke* Masayuki Hirukawa†
TH Köln Ryukoku University

4th April 2024

S.1 Introduction

In the proofs of lemmata below, equation numbers starting from “A” correspond to those in the Appendix. Before proceeding, we reproduce Appendix A.1 and A.2 as S.1.1 and S.1.2, respectively, for convenience.

S.2 Useful Formulae on the Gamma Function

Stirling’s Formula.

$$\Gamma(a+1) = \sqrt{2\pi} a^{a+1/2} \exp(-a) \left\{ 1 + \frac{1}{12a} + O(a^{-2}) \right\} \text{ as } a \rightarrow \infty. \quad (\text{S1})$$

Recursive Formula for the Lower Incomplete Gamma Function.

$$\gamma(a+1, z) = a\gamma(a, z) - z^a \exp(-z) \text{ for } a, z > 0. \quad (\text{S2})$$

Recursive Formula for the Polygamma Function.

$$\Psi^{(m)}(a+1) = \Psi^{(m)}(a) + \frac{(-1)^m m!}{a^{m+1}} \text{ for } a > 0 \text{ and } m \in \{0, 1, 2, \dots\}. \quad (\text{S3})$$

*Institute for Insurance Studies, TH Köln - University of Applied Sciences, Gustav-Heinemann-Ufer 54, 50968 Köln, Germany; e-mail: benedikt.funke@th-koeln.de.

†Faculty of Economics, Ryukoku University, 67 Tsukamoto-cho, Fukakusa, Fushimi-ku, Kyoto 612-8577, Japan; e-mail: hirukawa@econ.ryukoku.ac.jp.

S.3 Additional Notation

The following notation is adopted in the proofs: $R(a, z) = z^a \exp(-z) / \Gamma(a + 1)$ for $a, z > 0$; $\dot{K}_c^\pm(u) = \partial K_x^\pm(u) / \partial x|_{x=c}$; $\ddot{K}_c^\pm(u) = \partial^2 K_x^\pm(u) / \partial x^2|_{x=c}$; $\ddot{K}_c^\pm(u) = \partial^3 K_x^\pm(u) / \partial x^3|_{x=c}$; $H_i = \dot{K}_{t_0}^-(X_i) - \dot{K}_{t_0}^+(X_i)$; and $z^\pm = (t_0 \pm \Delta) / b$.

S.4 Proofs of Lemmata

S.4.1 Proof of Lemma A1

First two statements are direct outcomes from the proof of Lemma A.3 in Funke and Hirukawa (2023). The last statement can be obtained by equation (2.5) of Guo and Qi (2010) and (S1). ■

S.4.2 Proof of Lemma A2

The proof closely follows those for Lemmata 2 and 3 of Hirukawa et al. (2022). The case for $u = 0$ does not affect the derivation of the uniform bound, because the exponent on u in $K_x^\pm(u)$ is $a^\pm > 0$ so that $K_x^\pm(0) = \dot{K}_x^\pm(0) = 0$. For $u > 0$, we have

$$\dot{K}_x^\pm(u) = \frac{1}{b} \{ \ln u - \ln b - \Psi(a^\pm + 1) \} K_x^\pm(u).$$

Hence,

$$\left| \dot{K}_x^\pm(u) \right| \leq \frac{1}{b} \{ |\ln u| K_x^\pm(u) + |\ln b + \Psi(a^\pm + 1)| K_x^\pm(u) \}. \quad (\text{S4})$$

We work on the uniform bound of $|\ln b + \Psi(a^\pm + 1)| K_x^\pm(u)$ first. Observe that $|\ln b + \Psi(a^\pm + 1)| = |\ln x| + o(1)$, where the $o(1)$ approximation error is uniform on I_0 by Lemma A1. It is also the case that $|\ln x| \leq \max\{x^{-1}, x\}$, and we may take the $o(1)$ term no greater than 1 for a sufficiently large n . Then, $|\ln b + \Psi(a^\pm + 1)| \leq \max\{x^{-1}, x\} + 1$. It follows from $(x \pm \Delta)^{-1/2} = x^{-1/2} \{1 + o(1)\}$ and $o(1) \leq 1$ for a sufficiently large n in (A7) that

$$\begin{aligned} |\ln b + \Psi(a^\pm + 1)| K_x^\pm(u) &\leq \frac{b^{-1/2} \{1 + o(1)\}}{\sqrt{2\pi}} [\max\{x^{-3/2}, x^{1/2}\} + x^{-1/2}] \\ &\leq b^{-1/2} \sqrt{\frac{2}{\pi}} \left[\max\{\underline{t}^{-3/2}, \bar{t}^{1/2}\} + \underline{t}^{-1/2} \right] \end{aligned} \quad (\text{S5})$$

The relation $|\ln u| \leq \max\{u^{-1}, u\}$ again helps find the uniform bound of $|\ln u| K_x^\pm(u)$. Suppose that $|\ln u| \leq u^{-1}$. Because $u^{-1} K_x^\pm(u)$ is maximized at $u = x \pm \Delta - b$, it

holds that $|\ln u| K_x^\pm(u) \leq u^{-1} K_x^\pm(u) \leq (x \pm \Delta - b)^{-1} K_x^\pm(x \pm \Delta - b)$. By (S1),

$$\begin{aligned} & (x \pm \Delta - b)^{-1} K_x^\pm(x \pm \Delta - b) \\ &= \frac{b^{-1/2} e \{1 + o(1)\}}{\sqrt{2\pi}} \left(1 - \frac{1}{a^\pm}\right)^{a^\pm} \frac{1}{(x \pm \Delta - b) \sqrt{x \pm \Delta}}, \end{aligned}$$

where

$$\left(1 - \frac{1}{a^\pm}\right)^{a^\pm} = e^{-1} \{1 + o(1)\} \quad \text{and} \quad \frac{1}{(x \pm \Delta - b) \sqrt{x \pm \Delta}} = x^{-3/2} \{1 + o(1)\}.$$

It follows from $x^{-3/2} \leq \underline{t}^{-3/2}$ and $o(1) \leq 1$ for a sufficiently large n that

$$|\ln u| K_x^\pm(u) \leq \frac{b^{-1/2} \{1 + o(1)\}}{\sqrt{2\pi} x^{3/2}} \leq b^{-1/2} \sqrt{\frac{2}{\pi}} \underline{t}^{-3/2}.$$

Alternatively, suppose that $|\ln u| \leq u$. In this case, $u K_x^\pm(u)$ is maximized at $u = x \pm \Delta + b$, and thus $|\ln u| K_x^\pm(u) \leq u K_x^\pm(u) \leq (x \pm \Delta + b)^{-1} K_x^\pm(x \pm \Delta + b)$. Again by (S1),

$$\begin{aligned} & (x \pm \Delta + b)^{-1} K_x^\pm(x \pm \Delta + b) \\ &= \frac{b^{-1/2} e^{-1} \{1 + o(1)\}}{\sqrt{2\pi}} \left(1 + \frac{1}{a^\pm}\right)^{a^\pm} \frac{x \pm \Delta + b}{\sqrt{x \pm \Delta}}, \end{aligned}$$

where

$$\left(1 + \frac{1}{a^\pm}\right)^{a^\pm} = e \{1 + o(1)\} \quad \text{and} \quad \frac{x \pm \Delta + b}{\sqrt{x \pm \Delta}} = \sqrt{x} \{1 + o(1)\}.$$

Therefore,

$$|\ln u| K_x^\pm(u) \leq \frac{b^{-1/2} \sqrt{x} \{1 + o(1)\}}{\sqrt{2\pi}} \leq b^{-1/2} \sqrt{\frac{2}{\pi}} \bar{t}^{1/2},$$

by $x^{1/2} \leq \bar{t}^{1/2}$ and again $o(1) \leq 1$ for a sufficiently large n . Combining these two scenarios, we have

$$|\ln u| K_x^\pm(u) \leq \max \{u^{-1} K_x^\pm(u), u K_x^\pm(u)\} \leq b^{-1/2} \sqrt{\frac{2}{\pi}} \max \left\{ \underline{t}^{-3/2}, \bar{t}^{1/2} \right\}. \quad (\text{S6})$$

In the end, substituting (S5) and (S6) into (S4) yields

$$\sup_{(x,u) \in I_0 \times \mathbb{R}_+} \left| \dot{K}_x^\pm(u) \right| \leq b^{-3/2} \sqrt{\frac{2}{\pi}} \left[2 \max \left\{ \underline{t}^{-3/2}, \bar{t}^{1/2} \right\} + \underline{t}^{-1/2} \right] =: b^{-3/2} \sqrt{\frac{2}{\pi}} C_0.$$

This completes the proof. \blacksquare

S.4.3 Proof of Lemma A4

The lemma is proven by induction. The case for $m = 0$ is obvious by (S2) and $\Gamma(a+1) = a\Gamma(a)$. Next, suppose that the statement holds for some $m \geq 0$. Then,

$$\begin{aligned}
& \lambda^{m+1} \frac{\gamma(a+m+2, z)}{\Gamma(a+1)} \\
= & \lambda^{m+1} \frac{(a+m+1) \gamma(a+m+1, z) - z^{a+m+1} \exp(-z)}{\Gamma(a+1)} \\
= & \lambda(a+m+1) \{p_m P(a, z) - r_m R(a, z)\} - (\lambda z)^{m+1} R(a, z) \\
= & \{\lambda(a+m+1) p_m\} P(a, z) - \{\lambda(a+m+1) r_m + (\lambda z)^{m+1}\} R(a, z),
\end{aligned}$$

where the first and second equalities come from (S2) and the assumption of induction, respectively. The statement for $m+1$ is established by the correspondence of coefficients on $P(a, z)$ and $R(a, z)$. ■

S.4.4 Proof of Lemma A5

The first statement can be obtained by setting $x = t_0$ in (A12) and then substituting $\Phi(0) = 1/2$, $\phi(0) = 1/\sqrt{2\pi}$ and $\phi^{(1)}(0) = 0$. For the second statement, putting $x = t_0$ in (A18) gives

$$R(z^\pm, z_0) = \left\{ \frac{1 + O(1/z^\pm)}{\sqrt{2\pi}} \right\} \frac{1}{\sqrt{z^\pm}} \exp\{z^\pm (\ln \xi^\pm + 1 - \xi^\pm)\},$$

for $\xi^\pm := z_0/z^\pm = t_0/(t_0 \pm \Delta)$. When $x = t_0$, (A14) reduces to

$$\frac{1}{\sqrt{z^\pm}} = \frac{b^{1/2}}{\sqrt{t_0}} \left\{ 1 \mp \frac{\Delta}{2t_0} + O(\Delta^2) \right\}.$$

Moreover, $\exp\{z^\pm (\ln \xi^\pm + 1 - \xi^\pm)\}$ can be approximated by

$$\exp\{z^\pm (\ln \xi^\pm + 1 - \xi^\pm)\} = 1 - \frac{1}{2t_0} \left(\frac{\Delta^2}{b} \right) + O\left(\frac{\Delta^3}{b} \right).$$

Then, the second statement can be obtained by recognizing that $1 + O(1/z^\pm) = 1 + O(b)$. ■

S.4.5 Proof of Lemma A6

Observe that

$$E \left\{ \dot{K}_{t_0}^\pm(X_i) \right\} = \int_0^\infty \dot{K}_{t_0}^\pm(u) g(u) du + d_0 \int_0^{t_0} \dot{K}_{t_0}^\pm(u) du =: E_g^\pm + E_{d_0}^\pm,$$

where

$$\dot{K}_{t_0}^{\pm}(u) = \frac{1}{b} \left\{ \ln u - \ln b - \Psi(z^{\pm} + 1) \right\} K_{t_0}^{\pm}(u).$$

Then,

$$E \left\{ \hat{J}^{(1)}(t_0) \right\} = E \left\{ \dot{K}_{t_0}^{-}(X_i) \right\} - E \left\{ \dot{K}_{t_0}^{+}(X_i) \right\} = (E_g^{-} - E_g^{+}) + (E_{d_0}^{-} - E_{d_0}^{+}).$$

We start from approximating $(E_g^{-} - E_g^{+})$ and then proceed to $(E_{d_0}^{-} - E_{d_0}^{+})$. A minor modification of the proof for Theorem 2.1(i) of Funke and Hirukawa (2023) yields

$$E_g^{\pm} = g^{(1)}(t_0) \pm g^{(2)}(t_0) \Delta + O(b).$$

It follows that

$$E_g^{-} - E_g^{+} = -2g^{(2)}(t_0) \Delta + O(b).$$

Next, by Lemma A1 and a second-order Taylor expansion of $\ln u$ around $u = t_0$,

$$\begin{aligned} \ln u - \ln b - \Psi(z^{\pm} + 1) &= \ln u - \ln t_0 \mp \frac{\Delta}{t_0} - \frac{b}{2t_0} + O(b^2) \\ &\sim \left(\frac{u - t_0}{t_0} \right) - \frac{1}{2} \left(\frac{u - t_0}{t_0} \right)^2 \mp \frac{\Delta}{t_0} - \frac{b}{2t_0} + O(b^2). \end{aligned}$$

Then,

$$\begin{aligned} E_{d_0}^{\pm} &\sim \frac{d_0}{b} \left[\int_0^{t_0} \left(\frac{u - t_0}{t_0} \right) K_{t_0}^{\pm}(u) du - \frac{1}{2} \int_0^{t_0} \left(\frac{u - t_0}{t_0} \right)^2 K_{t_0}^{\pm}(u) du \right. \\ &\quad \left. + \left\{ \mp \frac{\Delta}{t_0} - \frac{b}{2t_0} + O(b^2) \right\} \int_0^{t_0} K_{t_0}^{\pm}(u) du \right]. \end{aligned}$$

Also observe that

$$\int_0^{t_0} u^m K_{t_0}^{\pm}(u) du =: b^m \frac{\gamma(z^{\pm} + m + 1, z_0)}{\Gamma(z^{\pm} + 1)}$$

for $m \geq 0$. Then, by Lemma A4 and straightforward calculations, we can obtain

$$\begin{aligned} \int_0^{t_0} K_{t_0}^{\pm}(u) du &= P(z^{\pm}, z_0) - R(z^{\pm}, z_0), \\ \int_0^{t_0} \left(\frac{u - t_0}{t_0} \right) K_{t_0}^{\pm}(u) du &= \left(\pm \frac{\Delta}{t_0} + \frac{b}{t_0} \right) P(z^{\pm}, z_0) - \left(1 \pm \frac{\Delta}{t_0} + \frac{b}{t_0} \right) R(z^{\pm}, z_0), \end{aligned}$$

and

$$\begin{aligned} &\int_0^{t_0} \left(\frac{u - t_0}{t_0} \right)^2 K_{t_0}^{\pm}(u) du \\ &= \left(\frac{b}{t_0} + \frac{2b^2}{t_0^2} \pm \frac{3\Delta b}{t_0^2} + \frac{\Delta^2}{t_0^2} \right) P(z^{\pm}, z_0) - \left(\frac{3b}{t_0} \pm \frac{\Delta}{t_0} + \frac{2b^2}{t_0^2} \pm \frac{3\Delta b}{t_0^2} + \frac{\Delta^2}{t_0^2} \right) R(z^{\pm}, z_0). \end{aligned}$$

It follows that

$$E_{d_0}^\pm \sim \frac{d_0}{b} \left[\left\{ \mp \frac{3\Delta b}{2t_0^2} + O(\Delta^2) \right\} P(z^\pm, z_0) - \left\{ 1 - \frac{b}{t_0} \mp \frac{\Delta}{2t_0} + O(\Delta^2) \right\} R(z^\pm, z_0) \right]$$

Substituting Lemma A5 and then making straightforward but tedious calculations, we have

$$E_{d_0}^- - E_{d_0}^+ = -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{1/2}} \right) + O\left(\frac{\Delta^3}{b^{3/2}} \right).$$

In the end,

$$\begin{aligned} E \left\{ \hat{J}^{(1)}(t_0) \right\} &= O(\Delta) + \left\{ -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{1/2}} \right) + O\left(\frac{\Delta^3}{b^{3/2}} \right) \right\} \\ &= -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{1/2}} \right) + O\left(\frac{\Delta^3}{b^{3/2}} \right) \\ &\sim -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{1/2}} \right). \end{aligned}$$

This completes the proof. ■

S.4.6 Proof of Lemma A7

Observe that

$$Var \left\{ \hat{J}^{(1)}(t_0) \right\} = \frac{1}{n} Var(H_i) = \frac{1}{n} \left\{ E(H_i^2) - E^2(H_i) \right\},$$

where $E(H_i) = E_{d_0}^- - E_{d_0}^+ = O(\Delta/b^{1/2})$ as shown above. Below we concentrate on approximating

$$E(H_i^2) = \int_0^\infty \left\{ \dot{K}_{t_0}^-(u) - \dot{K}_{t_0}^+(u) \right\}^2 f(u) du.$$

A difficulty arises at this stage because of asymmetry in $\dot{K}_x^-(\cdot) - \dot{K}_x^+(\cdot)$. If it were symmetric, we could first approximate the right-hand side by

$$\left\{ \frac{f(t_0^-) + f(t_0^+)}{2} \right\} \int_0^\infty \left\{ \dot{K}_{t_0}^-(u) - \dot{K}_{t_0}^+(u) \right\}^2 du.$$

However, it is not obvious whether this type of manipulation is still valid for the asymmetric case. Instead, our proof strategy is to decompose $f(\cdot)$ into two components so that $E(H_i^2) =: V_g + V_{d_0}$, where

$$\begin{aligned} V_g &= \int_0^\infty \left\{ \dot{K}_{t_0}^-(u) - \dot{K}_{t_0}^+(u) \right\}^2 g(u) du, \text{ and} \\ V_{d_0} &= d_0 \int_0^{t_0} \left\{ \dot{K}_{t_0}^-(u) - \dot{K}_{t_0}^+(u) \right\}^2 du. \end{aligned}$$

Below it is demonstrated that two integrals can be individually approximated by

$$V_g \sim g(t_0) \frac{3}{2\sqrt{\pi}t_0^{5/2}} \left(\frac{\Delta^2}{b^{5/2}} \right), \text{ and} \quad (\text{S7})$$

$$V_{d_0} \sim \left(\frac{d_0}{2} \right) \frac{3}{2\sqrt{\pi}t_0^{5/2}} \left(\frac{\Delta^2}{b^{5/2}} \right). \quad (\text{S8})$$

It should be recognized that (S7) and (S8) jointly establish the lemma, because $g(t_0) + d_0/2 = \{f(t_0^-) + f(t_0^+)\}/2$.

S.4.6.1 Proof of (S7)

Define $V_g^{2\pm} := \int_0^\infty \left\{ \dot{K}_{t_0}^\pm(u) \right\}^2 g(u) du$ and $V_g^{+-} := \int_0^\infty \dot{K}_{t_0}^-(u) \dot{K}_{t_0}^+(u) g(u) du$. Then, V_g can be decomposed into $V_g = V_g^{2-} + V_g^{2+} - 2V_g^{+-}$. In what follows, approximations to $V_g^{2\pm}$ and V_g^{+-} are derived separately. where $V_g^{2\pm} = \int_0^\infty \left\{ \dot{K}_{t_0}^\pm(u) \right\}^2 g(u) du$ and $V_g^{+-} = \int_0^\infty \dot{K}_{t_0}^-(u) \dot{K}_{t_0}^+(u) g(u) du$.

We start from finding an approximation to

$$V_g^{2\pm} = \frac{1}{b^2} \int_0^\infty \left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\}^2 \left\{ K_{t_0}^\pm(u) \right\}^2 g(u) du.$$

The integral can be further rewritten as $A^\pm(t_0) E \left[\left\{ \ln Y^\pm - \ln b - \Psi(z^\pm + 1) \right\}^2 g(Y^\pm) \right]$ for $Y^\pm \stackrel{d}{=} G(2z^\pm + 1, b/2)$ and $A^\pm(x)$ defined in (A8). By (S1), $A^\pm(t_0)$ can be approximated by

$$\begin{aligned} A^\pm(t_0) &= \frac{b^{-1}\Gamma(2z^\pm + 1)}{2^{2z^\pm+1}\Gamma^2(z^\pm + 1)} \\ &= \frac{b^{-1/2}}{2\sqrt{\pi}} (t_0 \pm \Delta)^{-1/2} \left\{ 1 - \frac{b}{8(t_0 \pm \Delta)} + O(b^2) \right\}. \end{aligned} \quad (\text{S9})$$

In addition,

$$\begin{aligned} &E \left[\left\{ \ln Y^\pm - \ln b - \Psi(z^\pm + 1) \right\}^2 g(Y^\pm) \right] \\ &= E \left\{ \ln Y^\pm - \ln b - \Psi(z^\pm + 1) \right\}^2 g(t_0) \{1 + o(1)\}. \end{aligned}$$

where, by the proof for Theorem 2.1(ii) of Funke and Hirukawa (2023) and Lemma A1,

$$\begin{aligned} &E \left\{ \ln Y^\pm - \ln b - \Psi(z^\pm + 1) \right\}^2 \\ &= \left\{ \ln 2 + \Psi(z^\pm + 1) - \Psi(2z^\pm + 1) \right\}^2 + \Psi^{(1)}(2z^\pm + 1) \\ &= \left\{ \frac{b}{4(t_0 \pm \Delta)} + O(b^2) \right\}^2 + \left\{ \frac{b}{2(t_0 \pm \Delta)} - \frac{b^2}{8(t_0 \pm \Delta)^2} + O(b^3) \right\} \\ &= \frac{b}{2(t_0 \pm \Delta)} \left\{ 1 - \frac{b}{8(t_0 \pm \Delta)} + O(b^2) \right\}. \end{aligned}$$

Hence,

$$\begin{aligned} V_g^{2\pm} &= \frac{b^{-3/2}g(t_0)\{1+o(1)\}}{4\sqrt{\pi}}(t_0\pm\Delta)^{-3/2}\left\{1-\frac{b}{8(t_0\pm\Delta)}+O(b^2)\right\}^2 \\ &= \frac{b^{-3/2}g(t_0)}{4\sqrt{\pi}t_0^{3/2}}\left\{1\mp\frac{3\Delta}{2t_0}-\frac{b}{4t_0}\pm\frac{\Delta b}{4t_0^2}+O(\Delta^2)\right\}. \end{aligned} \quad (\text{S10})$$

Next, we approximate

$$\begin{aligned} &V_g^{+-} \\ &= \frac{B(t_0)}{b^2}E\left[\{\ln Y_0 - \ln b - \Psi(z^- + 1)\}\{\ln Y_0 - \ln b - \Psi(z^+ + 1)\}g(Y_0)\right] \\ &= \frac{B(t_0)g(t_0)\{1+o(1)\}}{b^2}E\left[\{\ln Y_0 - \ln b - \Psi(z^- + 1)\}\{\ln Y_0 - \ln b - \Psi(z^+ + 1)\}\right] \end{aligned}$$

for $Y_0 \stackrel{d}{=} G(2z_0 + 1, b/2)$, where

$$B(t_0) := \frac{b^{-1}\Gamma(2z_0 + 1)}{2^{2z_0+1}\Gamma(z^- + 1)\Gamma(z^+ + 1)}.$$

Then, by (S1) and $b = o(\Delta^2/b)$,

$$\begin{aligned} &B(t_0) \\ &= \frac{b^{-1/2}}{2\sqrt{\pi}}\left(\frac{t_0}{t_0^2 - \Delta^2}\right)^{1/2}\left\{\left(\frac{t_0}{t_0 - \Delta}\right)^{t_0 - \Delta}\left(\frac{t_0}{t_0 + \Delta}\right)^{t_0 + \Delta}\right\}^{1/b}\left\{1 - \frac{b}{8t_0} + o(\Delta^2)\right\} \\ &= \frac{b^{-1/2}}{2\sqrt{\pi}\sqrt{t_0}}\{1 + O(\Delta^2)\}\left\{1 - \frac{1}{t_0}\left(\frac{\Delta^2}{b}\right) + o\left(\frac{\Delta^2}{b}\right)\right\}\{1 + O(b)\} \\ &= \frac{b^{-1/2}}{2\sqrt{\pi}\sqrt{t_0}}\left\{1 - \frac{1}{t_0}\left(\frac{\Delta^2}{b}\right) + o\left(\frac{\Delta^2}{b}\right)\right\}. \end{aligned} \quad (\text{S11})$$

It can be deduced again from by the proof for Theorem 2.1(ii) of Funke and Hirukawa (2023) and Lemma A1 that

$$\begin{aligned} &E\left[\{\ln Y_0 - \ln b - \Psi(z^- + 1)\}\{\ln Y_0 - \ln b - \Psi(z^+ + 1)\}\right] \\ &= \{\ln 2 + \Psi(z^- + 1) - \Psi(2z_0 + 1)\}\{\ln 2 + \Psi(z^+ + 1) - \Psi(2z_0 + 1)\} \\ &\quad + \Psi^{(1)}(2z_0 + 1) \\ &= \left\{-\frac{\Delta}{t_0} + O(b)\right\}\left\{\frac{\Delta}{t_0} + O(b)\right\} + \left\{\frac{b}{2t_0} + O(b^2)\right\} \\ &= \frac{b}{2t_0}\left\{1 - \frac{2}{t_0}\left(\frac{\Delta^2}{b}\right) + o\left(\frac{\Delta^2}{b}\right)\right\}. \end{aligned} \quad (\text{S12})$$

It follows from (S11) and (S12) that

$$V_g^{+-} = \frac{b^{-3/2}g(t_0)}{4\sqrt{\pi}t_0^{3/2}}\left\{1 - \frac{3}{t_0}\left(\frac{\Delta^2}{b}\right) + o\left(\frac{\Delta^2}{b}\right)\right\}. \quad (\text{S13})$$

In conclusion, by (S10), (S13) and $b = o(\Delta^2/b)$,

$$\begin{aligned} V_g &= (V_g^{2-} + V_g^{2+}) - 2V_g^{+-} \\ &= \frac{b^{-3/2}g(t_0)}{4\sqrt{\pi}t_0^{3/2}} \{2 + O(b)\} - \frac{b^{-3/2}g(t_0)}{4\sqrt{\pi}t_0^{3/2}} \left\{ 2 - \frac{6}{t_0} \left(\frac{\Delta^2}{b} \right) + o\left(\frac{\Delta^2}{b} \right) \right\} \\ &\sim g(t_0) \frac{3}{2\sqrt{\pi}t_0^{5/2}} \left(\frac{\Delta^2}{b^{5/2}} \right). \end{aligned}$$

S.4.6.2 Proof of (S8)

Denote $V_{d_0}^{2\pm} := d_0 \int_0^\infty \left\{ \dot{K}_{t_0}^\pm(u) \right\}^2 du$ and $V_{d_0}^{+-} := d_0 \int_0^\infty \dot{K}_{t_0}^-(u) \dot{K}_{t_0}^+(u) du$ so that $V_{d_0} = V_{d_0}^{2-} + V_{d_0}^{2+} - 2V_{d_0}^{+-}$. As above, $V_{d_0}^{2\pm}$ and $V_{d_0}^{+-}$ are approximated separately.

Below

$$V_{d_0}^{2\pm} = \frac{d_0 A^\pm(t_0)}{b^2} \int_0^{t_0} \left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\}^2 \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm + 1)} du$$

is approximated. Lemma A1 implies that

$$\begin{aligned} &\left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\}^2 \\ &= \left\{ \ln u - \ln t_0 \mp \frac{\Delta}{t_0} - \frac{b}{2t_0} + O(\Delta^2) \right\}^2 \\ &= \ln^2 u - 2 \left\{ \ln t_0 \pm \frac{\Delta}{t_0} + \frac{b}{2t_0} + O(\Delta^2) \right\} \ln u + \left\{ \ln t_0 \pm \frac{\Delta}{t_0} + \frac{b}{2t_0} + O(\Delta^2) \right\}^2. \end{aligned}$$

Combining this with a second-order Taylor expansion of $\ln^m u$ for $m \geq 1$ around $u = t_0$, i.e.,

$$\begin{aligned} \ln^m u &\approx \ln^m t_0 + m \ln^{m-1} t_0 \left(\frac{u - t_0}{t_0} \right) \\ &\quad + \frac{m}{2} \{ (m-1) \ln^{m-2} t_0 - \ln^{m-1} t_0 \} \left(\frac{u - t_0}{t_0} \right)^2, \end{aligned} \quad (\text{S14})$$

we have

$$\begin{aligned} &\left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\}^2 \\ &\sim \left\{ \mp \frac{2\Delta}{t_0} - \frac{b}{t_0} + O(\Delta^2) \right\} \left(\frac{u - t_0}{t_0} \right) + \left\{ 1 \pm \frac{\Delta}{t_0} + \frac{b}{2t_0} + O(\Delta^2) \right\} \left(\frac{u - t_0}{t_0} \right)^2 \\ &\quad + \left\{ \pm \frac{\Delta}{t_0} + \frac{b}{2t_0} + O(\Delta^2) \right\}^2. \end{aligned}$$

Observe that

$$\int_0^{t_0} u^m \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm + 1)} du = \left(\frac{b}{2} \right)^m \frac{\gamma(2z^\pm + m + 1, 2z_0)}{\Gamma(2z^\pm + 1)}$$

for $m \geq 0$. Then, Lemma A4 and straightforward calculations lead to

$$\begin{aligned}
& \int_0^{t_0} \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm+1)} du \\
&= P(2z^\pm, 2z_0) - R(2z^\pm, 2z_0), \\
& \int_0^{t_0} \left(\frac{u-t_0}{t_0}\right) \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm+1)} du \\
&= \left(\pm \frac{\Delta}{t_0} + \frac{b}{2t_0}\right) P(2z^\pm, 2z_0) - \left(1 \pm \frac{\Delta}{t_0} + \frac{b}{2t_0}\right) R(2z^\pm, 2z_0),
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^{t_0} \left(\frac{u-t_0}{t_0}\right)^2 \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm+1)} du \\
&= \left(\frac{b}{2t_0} + \frac{b^2}{2t_0^2} \pm \frac{3\Delta b}{2t_0^2} + \frac{\Delta^2}{t_0^2}\right) P(2z^\pm, 2z_0) \\
&\quad - \left(\frac{3b}{2t_0} \pm \frac{\Delta}{t_0} + \frac{b^2}{2t_0^2} \pm \frac{3\Delta b}{2t_0^2} + \frac{\Delta^2}{t_0^2}\right) R(2z^\pm, 2z_0).
\end{aligned}$$

It follows that

$$\begin{aligned}
& \int_0^{t_0} \left\{\ln u - \ln b - \Psi(z^\pm + 1)\right\}^2 \frac{u^{2z^\pm} \exp\{-u/(b/2)\}}{(b/2)^{2z^\pm+1} \Gamma(2z^\pm+1)} du \\
&\sim \left\{\frac{b}{2t_0} \pm \frac{\Delta b}{t_0^2} + O(\Delta^2)\right\} P(2z^\pm, 2z_0) \\
&\quad - \left\{\frac{b}{2t_0} \mp \frac{\Delta}{t_0} + O(\Delta^2)\right\} R(2z^\pm, 2z_0). \tag{S15}
\end{aligned}$$

Replacing b by $b/2$ in Lemma A5 also yields

$$P(2z^\pm, 2z_0) = \frac{1}{2} \mp \frac{1}{\sqrt{\pi}\sqrt{t_0}} \left(\frac{\Delta}{b^{1/2}}\right) + O\left(\frac{\Delta^2}{b^{1/2}}\right), \text{ and} \tag{S16}$$

$$R(2z^\pm, 2z_0) = \frac{b^{1/2}}{2\sqrt{\pi}\sqrt{t_0}} \left[1 \mp \frac{\Delta}{2t_0} - \frac{1}{t_0} \left(\frac{\Delta^2}{b}\right) + O\left\{\max\left(b, \frac{\Delta^3}{b}\right)\right\}\right]. \tag{S17}$$

Combining (S9), (S15), (S16), and (S17) and then recognizing that the $O(\Delta^2/b^{1/2})$ and $O\{\max(b, \Delta^3/b)\}$ terms are both $o(\Delta^2/b)$, we have

$$V_{d_0}^{2\pm} = d_0 \frac{b^{-3/2}}{4\sqrt{\pi}t_0^{3/2}} \left\{\frac{1}{2} \mp \frac{1}{\sqrt{\pi}\sqrt{t_0}} \left(\frac{\Delta}{b^{1/2}}\right) + o\left(\frac{\Delta}{b^{1/2}}\right)\right\}. \tag{S18}$$

Our next task is to find an approximation to

$$\begin{aligned}
V_{d_0}^{+-} &= d_0 \frac{B(t_0)}{b^2} \int_0^{t_0} \left[\{\ln u - \ln b - \Psi(z^- + 1)\} \{\ln u - \ln b - \Psi(z^+ + 1)\}\right. \\
&\quad \left. \times \frac{u^{2z_0} \exp(-2u/b)}{(b/2)^{2z_0+1} \Gamma(2z_0+1)}\right] du.
\end{aligned}$$

By Lemma A1,

$$2 \ln b + \Psi(z^- + 1) + \Psi(z^+ + 1) = 2 \ln t_0 + \frac{b}{t_0} - \frac{\Delta^2}{t_0^2} + o(\Delta^2),$$

and

$$\begin{aligned} & \{\ln b + \Psi(z^- + 1)\} \{\ln b + \Psi(z^+ + 1)\} \\ &= \ln^2 t_0 + \left(\frac{\ln t_0}{t_0}\right) b - \left(\frac{1 + \ln t_0}{t_0^2}\right) \Delta^2 + o(\Delta^2). \end{aligned}$$

Combining these with (S14) yields

$$\begin{aligned} & \{\ln u - \ln b - \Psi(z^- + 1)\} \{\ln u - \ln b - \Psi(z^+ + 1)\} \\ &= \ln^2 u - \{2 \ln b + \Psi(z^- + 1) + \Psi(z^+ + 1)\} \ln u \\ & \quad + \{\ln b + \Psi(z^- + 1)\} \{\ln b + \Psi(z^+ + 1)\} \\ & \sim \left\{ -\frac{b}{t_0} + \frac{\Delta^2}{t_0^2} + o(\Delta^2) \right\} \left(\frac{u - t_0}{t_0} \right) \\ & \quad + \left\{ 1 + \frac{b}{2t_0} - \frac{\Delta^2}{2t_0^2} + o(\Delta^2) \right\} \left(\frac{u - t_0}{t_0} \right)^2 - \frac{\Delta^2}{t_0^2} + o(\Delta^2). \end{aligned}$$

Notice that

$$\int_0^{t_0} u^m \frac{u^{2z_0} \exp\{-u/(b/2)\}}{(b/2)^{2z_0+1} \Gamma(2z_0+1)} du = \left(\frac{b}{2}\right)^m \frac{\gamma(2z_0+m+1, 2z_0)}{\Gamma(2z_0+1)}$$

for $m \geq 0$. Again by Lemma A4 and straightforward calculations, we have

$$\begin{aligned} \int_0^{t_0} \frac{u^{2z_0} \exp\{-u/(b/2)\}}{(b/2)^{2z_0+1} \Gamma(2z_0+1)} du &= P(2z_0, 2z_0) - R(2z_0, 2z_0), \\ \int_0^{t_0} \left(\frac{u - t_0}{t_0}\right) \frac{u^{2z_0} \exp\{-u/(b/2)\}}{(b/2)^{2z_0+1} \Gamma(2z_0+1)} du &= \frac{b}{2t_0} P(2z_0, 2z_0) - \left(1 + \frac{b}{2t_0}\right) R(2z_0, 2z_0), \end{aligned}$$

and

$$\begin{aligned} & \int_0^{t_0} \left(\frac{u - t_0}{t_0}\right)^2 \frac{u^{2z_0} \exp\{-u/(b/2)\}}{(b/2)^{2z_0+1} \Gamma(2z_0+1)} du \\ &= \left(\frac{b}{2t_0} + \frac{b^2}{2t_0^2}\right) P(2z_0, 2z_0) - \left(\frac{3b}{2t_0} + \frac{b^2}{2t_0^2}\right) R(2z_0, 2z_0). \end{aligned}$$

Hence,

$$\begin{aligned} & \int_0^{t_0} [\{\ln u - \ln b - \Psi(z^- + 1)\} \{\ln u - \ln b - \Psi(z^+ + 1)\}] \\ & \quad \times \frac{u^{2z_0} \exp\{-u/(b/2)\}}{(b/2)^{2z_0+1} \Gamma(2z_0+1)} du \\ & \sim \left\{ \frac{b}{2t_0} - \frac{\Delta^2}{t_0^2} + o(\Delta^2) \right\} P(2z_0, 2z_0) - \left\{ \frac{b}{2t_0} + o(\Delta^2) \right\} R(2z_0, 2z_0). \quad (\text{S19}) \end{aligned}$$

In addition, letting $x \downarrow 0$ in equation (1) of Pagurova (1965), we have

$$P(2z_0, 2z_0) = \frac{1}{2} + \frac{b^{1/2}}{6\sqrt{\pi}\sqrt{t_0}} + O(b^{3/2}). \quad (\text{S20})$$

Moreover, by (S1), $R(2z_0, 2z_0)$ can be approximated as

$$R(2z_0, 2z_0) = \frac{b^{1/2}}{2\sqrt{\pi}\sqrt{t_0}} \{1 + O(b)\}. \quad (\text{S21})$$

It follows from (S11), (S19), (S20), and (S21) that

$$V_{d_0}^{+-} = d_0 \frac{b^{-3/2}}{4\sqrt{\pi}t_0^{3/2}} \left\{ \frac{1}{2} - \frac{3}{2t_0} \left(\frac{\Delta^2}{b} \right) + o\left(\frac{\Delta^2}{b} \right) \right\}. \quad (\text{S22})$$

Substituting (S18) and (S22) into $V_{d_0} = (V_{d_0}^{2-} + V_{d_0}^{2+}) - 2V_{d_0}^{+-}$, we finally have

$$\begin{aligned} V_{d_0} &= d_0 \frac{b^{-3/2}}{4\sqrt{\pi}t_0^{3/2}} \left\{ 1 + o\left(\frac{\Delta}{b^{1/2}} \right) \right\} - d_0 \frac{b^{-3/2}}{4\sqrt{\pi}t_0^{3/2}} \left\{ 1 - \frac{3}{t_0} \left(\frac{\Delta^2}{b} \right) + o\left(\frac{\Delta^2}{b} \right) \right\} \\ &\sim \left(\frac{d_0}{2} \right) \frac{3}{2\sqrt{\pi}t_0^{5/2}} \left(\frac{\Delta^2}{b^{5/2}} \right). \end{aligned}$$

This completes the proof. \blacksquare

S.4.7 Proof of Lemma A8

It holds that

$$\begin{aligned} &\left| \left(\frac{b^{3/2}}{\Delta} \right) \hat{j}^{(2)}(\xi) - \left(-\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \right) \right| \\ &\leq \left(\frac{b^{3/2}}{\Delta} \right) \left| \hat{j}^{(2)}(\xi) - \hat{j}^{(2)}(t_0) \right| + \left(\frac{b^{3/2}}{\Delta} \right) \left| \hat{j}^{(2)}(t_0) - E \left\{ \hat{j}^{(2)}(t_0) \right\} \right| \\ &\quad + \left| \left(\frac{b^{3/2}}{\Delta} \right) E \left\{ \hat{j}^{(2)}(t_0) \right\} - \left(-\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \right) \right| \\ &= D_1 + D_2 + D_3 \quad (\text{say}). \end{aligned}$$

Then, the proof boils down to establishing the following three statements:

$$D_1 = \left(\frac{b^{3/2}}{\Delta} \right) \left| \hat{j}^{(2)}(\xi) - \hat{j}^{(2)}(t_0) \right| = o_p(1). \quad (\text{S23})$$

$$D_2 = \left(\frac{b^{3/2}}{\Delta} \right) \left| \hat{j}^{(2)}(t_0) - E \left\{ \hat{j}^{(2)}(t_0) \right\} \right| = o_p(1). \quad (\text{S24})$$

$$D_3 = \left| \left(\frac{b^{3/2}}{\Delta} \right) E \left\{ \hat{j}^{(2)}(t_0) \right\} - \left(-\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \right) \right| = o(1). \quad (\text{S25})$$

We work on (S25) first and then proceed to (S23) and (S24).

S.4.7.1 Proof of (S25)

The proof is similar to that of Lemma A6, and thus we provide only its outline. Because

$$E \left\{ \ddot{K}_{t_0}^{\pm}(X_i) \right\} = \int_0^{\infty} \dot{K}_{t_0}^{\pm}(u) g(u) du + d_0 \int_0^{t_0} \dot{K}_{t_0}^{\pm}(u) du =: I_g^{\pm} + I_{d_0}^{\pm},$$

where

$$\ddot{K}_{t_0}^{\pm}(u) = \frac{1}{b^2} \left[\left\{ \ln u - \ln b - \Psi(z^{\pm} + 1) \right\}^2 - \Psi^{(1)}(z^{\pm} + 1) \right] K_{t_0}^{\pm}(u),$$

it holds that

$$E \left\{ \hat{J}^{(2)}(t_0) \right\} = E \left\{ \ddot{K}_{t_0}^{-}(X_i) \right\} - E \left\{ \ddot{K}_{t_0}^{+}(X_i) \right\} = (I_g^{-} - I_g^{+}) + (I_{d_0}^{-} - I_{d_0}^{+}).$$

A similar procedure to the proof of Lemma A6 yields

$$I_g^{\pm} = g^{(2)}(t_0) \pm g^{(3)}(t_0) \Delta + o(\Delta)$$

so that

$$I_g^{-} - I_g^{+} = -2g^{(3)}(t_0) \Delta + o(\Delta).$$

It can be also found via straightforward but tedious calculations that

$$I_{d_0}^{\pm} \sim \frac{d_0}{b^2} \left[\left\{ -\frac{\Delta^2}{2t_0^2} \pm \frac{3\Delta b}{2t_0^2} + o(\Delta b) \right\} P(z^{\pm}, z_0) - \left\{ \mp \frac{\Delta}{t_0} + \frac{b}{t_0} + o(b) \right\} R(z^{\pm}, z_0) \right].$$

Then, using Lemma A5 yields

$$\begin{aligned} I_{d_0}^{-} - I_{d_0}^{+} &= \frac{d_0}{b^2} \left[\left\{ O(\Delta b) + O\left(\frac{\Delta^3}{b^{1/2}}\right) \right\} - \frac{b^{1/2}}{\sqrt{2\pi}\sqrt{t_0}} \left\{ \frac{2\Delta}{t_0} + O(\Delta b) \right\} \right] \\ &= -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{3/2}} \right) + O\left(\frac{\Delta^3}{b^{1/2}}\right). \end{aligned}$$

Therefore,

$$\begin{aligned} E \left\{ \hat{J}^{(2)}(t_0) \right\} &= O(\Delta) + \left\{ -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{3/2}} \right) + O\left(\frac{\Delta^3}{b^{1/2}}\right) \right\} \\ &= -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{3/2}} \right) + O(\Delta) \\ &\sim -\sqrt{\frac{2}{\pi}} \frac{d_0}{t_0^{3/2}} \left(\frac{\Delta}{b^{3/2}} \right), \end{aligned}$$

and thus (S25) is demonstrated.

S.4.7.2 Proof of (S23)

By a first-order Taylor expansion of $\hat{J}^{(2)}(\xi)$ around $\xi = t_0$,

$$\begin{aligned}\hat{J}^{(2)}(\xi) - \hat{J}^{(2)}(t_0) &= \hat{J}^{(3)}(\tau)(\xi - t_0) \\ &= \hat{J}^{(3)}(t_0)(t_0)(\xi - t_0) + \left\{ \hat{J}^{(3)}(\tau) - \hat{J}^{(3)}(t_0) \right\}(\xi - t_0)\end{aligned}$$

for some τ between \tilde{t} and t_0 . It can be shown that the second term is of smaller order than the first term by Lipschitz continuity of $\hat{J}^{(3)}(\cdot)$. To put it in another way, the order of magnitude in $(b^{3/2}/\Delta) \left| \hat{J}^{(2)}(\xi) - \hat{J}^{(2)}(t_0) \right|$ is determined by that of $(b^{3/2}/\Delta) \left| \hat{J}^{(3)}(t_0) \right| |\xi - t_0|$.

In what follows, the order of magnitude in $\hat{J}^{(3)}(t_0)$ can be found via the identity $\hat{J}^{(3)}(t_0) = E \left\{ \hat{J}^{(3)}(t_0) \right\} + \left[\hat{J}^{(3)}(t_0) - E \left\{ \hat{J}^{(3)}(t_0) \right\} \right]$. As in the proofs of Lemma A6 and (S25),

$$\begin{aligned}E \left\{ \hat{J}^{(3)}(t_0) \right\} &= E \left\{ \ddot{K}_{t_0}^-(X_i) \right\} - E \left\{ \ddot{K}_{t_0}^+(X_i) \right\} \\ &= \left\{ \int_0^{t_0} \ddot{K}_{t_0}^-(u) g(u) du - \int_0^{t_0} \ddot{K}_{t_0}^+(u) g(u) du \right\} \\ &\quad + d_0 \left\{ \int_0^{t_0} \ddot{K}_{t_0}^-(u) du - \int_0^{t_0} \ddot{K}_{t_0}^+(u) du \right\},\end{aligned}$$

where

$$\begin{aligned}\ddot{K}_{t_0}^\pm(u) &= \frac{1}{b^3} \left[\left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\}^3 \right. \\ &\quad \left. - 3\Psi^{(1)}(z^\pm + 1) \left\{ \ln u - \ln b - \Psi(z^\pm + 1) \right\} - \Psi^{(2)}(z^\pm + 1) \right] K_{t_0}^\pm(u),\end{aligned}$$

and $\left| \int_0^{t_0} \ddot{K}_{t_0}^-(u) g(u) du - \int_0^{t_0} \ddot{K}_{t_0}^+(u) g(u) du \right|$ is at most $O(1)$ by Lemma A1 and boundedness of $g^{(3)}(\cdot)$ on I_0 . Below it turns out that $\left| d_0 \left\{ \int_0^{t_0} \ddot{K}_{t_0}^-(u) du - \int_0^{t_0} \ddot{K}_{t_0}^+(u) du \right\} \right|$ is the dominant term in $\left| E \left\{ \hat{J}^{(3)}(t_0) \right\} \right|$. A first-order Taylor expansion of $\ddot{K}_{t_0}^\pm(u)$ around $\Delta = 0$ implies that

$$\begin{aligned}\ddot{K}_{t_0}^-(u) - \ddot{K}_{t_0}^+(u) &\sim 2 \left(\frac{\Delta}{b^4} \right) \left[3 \left\{ \ln u - \ln b - \Psi(z_0 + 1) \right\}^2 \Psi^{(1)}(z_0 + 1) \right. \\ &\quad \left. + 3 \left\{ \ln u - \ln b - \Psi(z_0 + 1) \right\} \Psi^{(2)}(z_0 + 1) \right. \\ &\quad \left. - 3 \left\{ \Psi^{(1)}(z_0 + 1) \right\}^2 + \Psi^{(3)}(z_0 + 1) \right] K_{t_0}(u).\end{aligned}$$

Then, by Lemma A1 and a similar procedure to the proof of Lemma A6, it can be found that

$$\left| \int_0^{t_0} \ddot{K}_{t_0}^-(u) du - \int_0^{t_0} \ddot{K}_{t_0}^+(u) du \right| = O \left(\frac{\Delta}{b^4} \right) O(b^2) = O \left(\frac{\Delta}{b^2} \right).$$

Therefore, $\left| E \left\{ \hat{J}^{(3)}(t_0) \right\} \right| \leq O(1) + O(\Delta/b^2) = O(\Delta/b^2)$.

Furthermore,

$$Var \left\{ \hat{J}^{(3)}(t_0) \right\} = \frac{1}{n} \left[E \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\}^2 - E^2 \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\} \right],$$

where $\left| E \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\} \right| = \left| E \left\{ \hat{J}^{(3)}(t_0) \right\} \right| = O(\Delta/b^2)$ as above. By Lemma A1 and a similar procedure to the proof of Lemma A7, it can be shown that

$$E \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\}^2 = O\left(\frac{\Delta^2}{b^8}\right) O(b^{7/2}) = O\left(\frac{\Delta^2}{b^{9/2}}\right).$$

It follows that $Var \left\{ \hat{J}^{(3)}(t_0) \right\} = O\left\{ \Delta^2 / (nb^{9/2}) \right\}$.

In conclusion,

$$\hat{J}^{(3)}(t_0) = O\left(\frac{\Delta}{b^2}\right) + O_p\left(\sqrt{\frac{\Delta^2}{nb^{9/2}}}\right).$$

Because ξ can be expressed as $\xi = \theta \hat{t} + (1 - \theta)t_0$ for some $\theta \in [0, 1]$ and $|\hat{t} - t_0| = o_p(b^{1/2})$ by assumption, it holds that $|\xi - t_0| \leq |\hat{t} - t_0| = o_p(b^{1/2})$. Therefore,

$$\begin{aligned} \left(\frac{b^{3/2}}{\Delta}\right) \left| \hat{J}^{(3)}(t_0) \right| |\xi - t_0| &= \left(\frac{b^{3/2}}{\Delta}\right) \left\{ O\left(\frac{\Delta}{b^2}\right) + O_p\left(\sqrt{\frac{\Delta^2}{nb^{9/2}}}\right) \right\} o_p(b^{1/2}) \\ &= o_p(1) + o_p\left(\frac{1}{\sqrt{nb^{1/2}}}\right) \xrightarrow{p} 0, \end{aligned}$$

which establishes (S23).

S.4.7.3 Proof of (S24)

It suffices to show that $Var \left\{ (b^{3/2}/\Delta) \hat{J}^{(2)}(t_0) \right\} = o(1)$. To do so, again we focus on the order of magnitude in $E \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\}^2$. A first-order Taylor expansion of $\ddot{K}_{t_0}^\pm(u)$ around $\Delta = 0$ implies that

$$\begin{aligned} &\ddot{K}_{t_0}^-(u) - \ddot{K}_{t_0}^+(u) \\ &\sim 2 \left(\frac{\Delta}{b^3}\right) [-\{\ln u - \ln b - \Psi(z_0 + 1)\}]^3 \\ &\quad + 3 \{\ln u - \ln b - \Psi(z_0 + 1)\} \Psi^{(1)}(z_0 + 1) + \Psi^{(2)}(z_0 + 1) \left] K_{t_0}(u). \end{aligned}$$

Then, by a similar procedure to the proof of (S23) above,

$$E \left\{ \ddot{K}_{t_0}^-(X_i) - \ddot{K}_{t_0}^+(X_i) \right\}^2 = O\left(\frac{\Delta^2}{b^6}\right) O(b^{5/2}) = O\left(\frac{\Delta^2}{b^{7/2}}\right).$$

It follows that

$$Var \left\{ \left(\frac{b^{3/2}}{\Delta}\right) \hat{J}^{(2)}(t_0) \right\} = \left(\frac{b^3}{\Delta^2}\right) O\left(\frac{\Delta^2}{nb^{7/2}}\right) = O\left(\frac{1}{nb^{1/2}}\right) \rightarrow 0.$$

This completes the proof. \blacksquare

S.4.8 Proof of Lemma A9

Basically, a similar strategy to the proofs of (S23) and (S24) in Lemma A8 may be taken. The derivation is straightforward but much more tedious, and thus details are omitted. ■

References

- [1] Funke, B., and M. Hirukawa (2023): “Density Derivative Estimation Using Asymmetric Kernels,” *Journal of Nonparametric Statistics*, forthcoming.
- [2] Guo, B.-N., and F. Qi (2010): “Some Properties of the Psi and Polygamma Functions,” *Hacettepe Journal of Mathematics and Statistics*, 39, 219-231.
- [3] Hirukawa, M., Murtazashvili I., and A. Prokhorov (2022): “Uniform Convergence Rates for Nonparametric Estimators Smoothed by the Beta Kernel,” *Scandinavian Journal of Statistics*, 49, 1353-1382.
- [4] Pagurova, V. I. (1965): “An Asymptotic Formula for the Incomplete Gamma Function,” *USSR Computational Mathematics and Mathematical Physics*, 5, 162-166.